

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
7 June 2001 (07.06.2001)

PCT

(10) International Publication Number
WO 01/40466 A2

(51) International Patent Classification ⁷ :	C12N 15/12, C07K 14/47, 14/705, C12N 15/62, C07K 16/18, 16/28, G01N 33/53, A61K 38/17, C12Q 1/68	PCT/US99/30999	20 December 1999 (20.12.1999)	US
		PCT/US99/31243	30 December 1999 (30.12.1999)	US
(21) International Application Number:	PCT/US00/32678	PCT/US99/31274	30 December 1999 (30.12.1999)	US
(22) International Filing Date:	1 December 2000 (01.12.2000)	PCT/US00/00219	5 January 2000 (05.01.2000)	US
		PCT/US00/00277	6 January 2000 (06.01.2000)	US
(25) Filing Language:	English	PCT/US00/00376	6 January 2000 (06.01.2000)	US
		PCT/US00/03565	11 February 2000 (11.02.2000)	US
(26) Publication Language:	English	PCT/US00/04341	18 February 2000 (18.02.2000)	US
		PCT/US00/04342	18 February 2000 (18.02.2000)	US
(30) Priority Data:	PCT/US99/28301	PCT/US00/04414	22 February 2000 (22.02.2000)	US
	1 December 1999 (01.12.1999)	PCT/US00/04914	24 February 2000 (24.02.2000)	US
	PCT/US99/28634	PCT/US00/05004	24 February 2000 (24.02.2000)	US
	1 December 1999 (01.12.1999)	PCT/US00/05601	1 March 2000 (01.03.2000)	US
	PCT/US99/28551	PCT/US00/05841	2 March 2000 (02.03.2000)	US
	2 December 1999 (02.12.1999)	60/187,202	3 March 2000 (03.03.2000)	US
	PCT/US99/28564	PCT/US00/06319	10 March 2000 (10.03.2000)	US
	2 December 1999 (02.12.1999)	PCT/US00/06884	15 March 2000 (15.03.2000)	US
	PCT/US99/28565	PCT/US00/07377	20 March 2000 (20.03.2000)	US
	2 December 1999 (02.12.1999)	PCT/US00/07532	21 March 2000 (21.03.2000)	US
	60/170,262			
	9 December 1999 (09.12.1999)			
	PCT/US99/30095			
	16 December 1999 (16.12.1999)			
	PCT/US99/30911			
	20 December 1999 (20.12.1999)			

[Continued on next page]

(54) Title: SECRETED AND TRANSMEMBRANE POLYPEPTIDES AND NUCLEIC ACIDS ENCODING THE SAME

MSTMFADTLLIVFISVCTALLAEGITWVLVYRTDKYKRLKAEVEKQSKKLEKKKETITESAGR
 QQKKKIERQEEKLKNNNRDLMSVRMKSMAFGFCFTALMGMFNSIFDGRVVAKLFPFTLSYIQ
 GLSHRNLLGDDTTDCSFIFLYILCTMSIRQNIQKILGLAPSRATKQAGGFLGPPPPSGKFS

Important features:**Signal peptide:**

amino acids 1-22

N-myristoylation sites.

amino acids 103-109, 163-169

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 53-57

(57) Abstract: The present invention is directed to novel polypeptides and to nucleic acid molecules encoding those polypeptides. Also provided herein are vectors and host cells comprising those nucleic acid sequences, chimeric polypeptide molecules comprising the polypeptides of the present invention fused to heterologous polypeptide sequences, antibodies which bind to the polypeptides of the present invention and to methods for producing the polypeptides of the present invention.



WO 01/40466 A2



- | | | | |
|----------------|--------------------------------|----|--|
| PCT/US00/08439 | 30 March 2000 (30.03.2000) | US | Los Altos, CA 94024 (US). SMITH, Victoria [AU/US]; |
| PCT/US00/13705 | 17 May 2000 (17.05.2000) | US | 19 Dwight Road, Burlingame, CA 94010 (US). STEW- |
| PCT/US00/14042 | 22 May 2000 (22.05.2000) | US | ART, Timothy, A. [US/US]; 465 Douglass Street, San |
| PCT/US00/14941 | 30 May 2000 (30.05.2000) | US | Francisco, CA 94114 (US). TUMAS, Daniel [US/US]; |
| PCT/US00/15264 | 2 June 2000 (02.06.2000) | US | 3 Rae Avenue, Orinda, CA 94563 (US). WATANABE, |
| 60/209,832 | 5 June 2000 (05.06.2000) | US | Colin, K. [US/US]; 128 Corliss Drive, Moraga, CA 94556 |
| PCT/US00/20710 | 28 July 2000 (28.07.2000) | US | (US). WOOD, William, I. [US/US]; 35 Southdown Court, |
| PCT/US00/22031 | 11 August 2000 (11.08.2000) | US | Hillsborough, CA 94010 (US). ZHANG, Zemin [CN/US]; |
| PCT/US00/23522 | 23 August 2000 (23.08.2000) | US | 876 Taurus Drive, Foster City, CA 94404 (US). |
| PCT/US00/23328 | 24 August 2000 (24.08.2000) | US | |
| 60/000,000 | 15 September 2000 (15.09.2000) | US | (74) Agents: KRESNAK, Mark, T. et al.; Genentech, Inc., |
| PCT/US00/30952 | 8 November 2000 (08.11.2000) | US | MS49, 1 DNA Way, South San Francisco, CA 94080-4990 |
| PCT/US00/30873 | 10 November 2000 (10.11.2000) | US | (US). |
- (71) **Applicant (for all designated States except US): GENEN-TECH, INC.** [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only): BAKER, Kevin, P.** [GB/US]; 14006 Indian Run Drive, Darnestown, MD 20878 (US). **BERESINI, Maureen** [US/US]; 611 Stetson Street, Moss Beach, CA 94038 (US). **DEFORGE, Laura** [US/US]; 1175 Manzanita Drive, Pacifica, CA 94044 (US). **DESNOYERS, Luc** [CA/US]; 2050 Stockton Street, San Francisco, CA 94133 (US). **FILVAROFF, Ellen** [US/US]; 538 18th Avenue, San Francisco, CA 94121 (US). **GAO, Wei-Qiang** [CN/US]; 641 Pilgrim Drive, Foster City, CA 94404 (US). **GERRITSEN, Mary, E.** [CA/US]; 541 Parrott Drive, San Mateo, CA 94402 (US). **GODDARD, Audrey** [CA/US]; 110 Congo Street, San Francisco, CA 94131 (US). **GODOWSKI, Paul, J.** [US/US]; 2627 Easton Drive, Burlingame, CA 94010 (US). **GURNEY, Austin, L.** [US/US]; 1 Debbie Lane, Belmont, CA 94002 (US). **SHERWOOD, Steven** [US/US]; 995 Lundy Lane,
- (81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
 — Without international search report and to be republished upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

SECRETED AND TRANSMEMBRANE POLYPEPTIDES AND NUCLEIC ACIDS ENCODING THE SAME

FIELD OF THE INVENTION

The present invention relates generally to the identification and isolation of novel DNA and to the
5 recombinant production of novel polypeptides.

BACKGROUND OF THE INVENTION

Extracellular proteins play important roles in, among other things, the formation, differentiation and
maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration,
10 differentiation, or interaction with other cells, is typically governed by information received from other cells
and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance,
mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which
are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. These secreted
polypeptides or signaling molecules normally pass through the cellular secretory pathway to reach their site of
15 action in the extracellular environment.

Secreted proteins have various industrial applications, including as pharmaceuticals, diagnostics,
biosensors and bioreactors. Most protein drugs available at present, such as thrombolytic agents, interferons,
interleukins, erythropoietins, colony stimulating factors, and various other cytokines, are secretory proteins.
Their receptors, which are membrane proteins, also have potential as therapeutic or diagnostic agents. Efforts
20 are being undertaken by both industry and academia to identify new, native secreted proteins. Many efforts are
focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel
secreted proteins. Examples of screening methods and techniques are described in the literature [see, for
example, Klein et al., Proc. Natl. Acad. Sci. 93:7108-7113 (1996); U.S. Patent No. 5,536,637].

Membrane-bound proteins and receptors can play important roles in, among other things, the formation,
25 differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation,
migration, differentiation, or interaction with other cells, is typically governed by information received from
other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides
(for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and
hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins.
30 Such membrane-bound proteins and cell receptors include, but are not limited to, cytokine receptors, receptor
kinases, receptor phosphatases, receptors involved in cell-cell interactions, and cellular adhesion molecules like
selectins and integrins. For instance, transduction of signals that regulate cell growth and differentiation is
regulated in part by phosphorylation of various cellular proteins. Protein tyrosine kinases, enzymes that catalyze
that process, can also act as growth factor receptors. Examples include fibroblast growth factor receptor and

nerve growth factor receptor.

Membrane-bound proteins and receptor molecules have various industrial applications, including as pharmaceutical and diagnostic agents. Receptor immunoadhesins, for instance, can be employed as therapeutic agents to block receptor-ligand interactions. The membrane-bound proteins can also be employed for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction.

Efforts are being undertaken by both industry and academia to identify new, native receptor or membrane-bound proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel receptor or membrane-bound proteins.

SUMMARY OF THE INVENTION

In one embodiment, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a PRO polypeptide.

In one aspect, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule encoding a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

In other aspects, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94%

nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule comprising the coding sequence of a full-length PRO polypeptide cDNA as disclosed herein, the coding sequence of a PRO polypeptide lacking the signal peptide as disclosed herein, the coding sequence of an extracellular domain of a transmembrane PRO polypeptide, with or without the signal peptide, as disclosed herein or the coding sequence of any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

In a further aspect, the invention concerns an isolated nucleic acid molecule comprising a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule that encodes the same mature polypeptide encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein, or (b) the complement of the DNA molecule of (a).

Another aspect the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding a PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated, or is complementary to such encoding nucleotide sequence, wherein the transmembrane domain(s) of such polypeptide are disclosed herein. Therefore, soluble extracellular domains of the herein described PRO polypeptides are contemplated.

Another embodiment is directed to fragments of a PRO polypeptide coding sequence, or the complement thereof, that may find use as, for example, hybridization probes, for encoding fragments of a PRO polypeptide that may optionally encode a polypeptide comprising a binding site for an anti-PRO antibody or as antisense oligonucleotide probes. Such nucleic acid fragments are usually at least about 10 nucleotides in length, alternatively at least about 15 nucleotides in length, alternatively at least about 20 nucleotides in length, alternatively at least about 30 nucleotides in length, alternatively at least about 40 nucleotides in length, alternatively at least about 50 nucleotides in length, alternatively at least about 60 nucleotides in length, alternatively at least about 70 nucleotides in length, alternatively at least about 80 nucleotides in length, alternatively at least about 90 nucleotides in length, alternatively at least about 100 nucleotides in length, alternatively at least about 110 nucleotides in length, alternatively at least about 120 nucleotides in length,

alternatively at least about 130 nucleotides in length, alternatively at least about 140 nucleotides in length, alternatively at least about 150 nucleotides in length, alternatively at least about 160 nucleotides in length, alternatively at least about 170 nucleotides in length, alternatively at least about 180 nucleotides in length, alternatively at least about 190 nucleotides in length, alternatively at least about 200 nucleotides in length, alternatively at least about 250 nucleotides in length, alternatively at least about 300 nucleotides in length, alternatively at least about 350 nucleotides in length, alternatively at least about 400 nucleotides in length, alternatively at least about 450 nucleotides in length, alternatively at least about 500 nucleotides in length, alternatively at least about 600 nucleotides in length, alternatively at least about 700 nucleotides in length, alternatively at least about 800 nucleotides in length, alternatively at least about 900 nucleotides in length and alternatively at least about 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length. It is noted that novel fragments of a PRO polypeptide-encoding nucleotide sequence may be determined in a routine manner by aligning the PRO polypeptide-encoding nucleotide sequence with other known nucleotide sequences using any of a number of well known sequence alignment programs and determining which PRO polypeptide-encoding nucleotide sequence fragment(s) are novel. All of such PRO polypeptide-encoding nucleotide sequences are contemplated herein. Also contemplated are the PRO polypeptide fragments encoded by these nucleotide molecule fragments, preferably those PRO polypeptide fragments that comprise a binding site for an anti-PRO antibody.

In another embodiment, the invention provides isolated PRO polypeptide encoded by any of the isolated nucleic acid sequences hereinabove identified.

In a certain aspect, the invention concerns an isolated PRO polypeptide, comprising an amino acid sequence having at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein.

In a further aspect, the invention concerns an isolated PRO polypeptide comprising an amino acid sequence having at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid

sequence identity, alternatively at least about 82 % amino acid sequence identity, alternatively at least about 83 % amino acid sequence identity, alternatively at least about 84 % amino acid sequence identity, alternatively at least about 85 % amino acid sequence identity, alternatively at least about 86 % amino acid sequence identity, alternatively at least about 87 % amino acid sequence identity, alternatively at least about 88 % amino acid sequence identity, alternatively at least about 89 % amino acid sequence identity, alternatively at least about 90 % amino acid sequence identity, alternatively at least about 91 % amino acid sequence identity, alternatively at least about 92 % amino acid sequence identity, alternatively at least about 93 % amino acid sequence identity, alternatively at least about 94 % amino acid sequence identity, alternatively at least about 95 % amino acid sequence identity, alternatively at least about 96 % amino acid sequence identity, alternatively at least about 97 % amino acid sequence identity, alternatively at least about 98 % amino acid sequence identity and alternatively at least about 99 % amino acid sequence identity to an amino acid sequence encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein.

In a specific aspect, the invention provides an isolated PRO polypeptide without the N-terminal signal sequence and/or the initiating methionine and is encoded by a nucleotide sequence that encodes such an amino acid sequence as hereinbefore described. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

Another aspect the invention provides an isolated PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

In yet another embodiment, the invention concerns agonists and antagonists of a native PRO polypeptide as defined herein. In a particular embodiment, the agonist or antagonist is an anti-PRO antibody or a small molecule.

In a further embodiment, the invention concerns a method of identifying agonists or antagonists to a PRO polypeptide which comprise contacting the PRO polypeptide with a candidate molecule and monitoring a biological activity mediated by said PRO polypeptide. Preferably, the PRO polypeptide is a native PRO polypeptide.

In a still further embodiment, the invention concerns a composition of matter comprising a PRO polypeptide, or an agonist or antagonist of a PRO polypeptide as herein described, or an anti-PRO antibody, in combination with a carrier. Optionally, the carrier is a pharmaceutically acceptable carrier.

Another embodiment of the present invention is directed to the use of a PRO polypeptide, or an agonist or antagonist thereof as hereinbefore described, or an anti-PRO antibody, for the preparation of a medicament useful in the treatment of a condition which is responsive to the PRO polypeptide, an agonist or antagonist thereof or an anti-PRO antibody.

In other embodiments of the present invention, the invention provides vectors comprising DNA encoding any of the herein described polypeptides. Host cell comprising any such vector are also provided. By way of example, the host cells may be CHO cells, *E. coli*, or yeast. A process for producing any of the herein described polypeptides is further provided and comprises culturing host cells under conditions suitable for expression of the desired polypeptide and recovering the desired polypeptide from the cell culture.

5 In other embodiments, the invention provides chimeric molecules comprising any of the herein described polypeptides fused to a heterologous polypeptide or amino acid sequence. Example of such chimeric molecules comprise any of the herein described polypeptides fused to an epitope tag sequence or a Fc region of an immunoglobulin.

10 In another embodiment, the invention provides an antibody which binds, preferably specifically, to any of the above or below described polypeptides. Optionally, the antibody is a monoclonal antibody, humanized antibody, antibody fragment or single-chain antibody.

In yet other embodiments, the invention provides oligonucleotide probes which may be useful for isolating genomic and cDNA nucleotide sequences, measuring or detecting expression of an associated gene or as antisense probes, wherein those probes may be derived from any of the above or below described nucleotide
15 sequences. Preferred probe lengths are described above.

In yet other embodiments, the present invention is directed to methods of using the PRO polypeptides of the present invention for a variety of uses based upon the functional biological assay data presented in the Examples below.

20 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a nucleotide sequence (SEQ ID NO:1) of a native sequence PRO177 cDNA, wherein SEQ ID NO:1 is a clone designated herein as "DNA16438-1387".

Figure 2 shows the amino acid sequence (SEQ ID NO:2) derived from the coding sequence of SEQ ID NO:1 shown in Figure 1.

25 Figure 3 shows a nucleotide sequence (SEQ ID NO:3) of a native sequence PRO3574 cDNA, wherein SEQ ID NO:3 is a clone designated herein as "DNA19360-2552".

Figure 4 shows the amino acid sequence (SEQ ID NO:4) derived from the coding sequence of SEQ ID NO:3 shown in Figure 3.

30 Figure 5 shows a nucleotide sequence (SEQ ID NO:5) of a native sequence PRO1280 cDNA, wherein SEQ ID NO:5 is a clone designated herein as "DNA33455-1548".

Figure 6 shows the amino acid sequence (SEQ ID NO:6) derived from the coding sequence of SEQ ID NO:5 shown in Figure 5.

Figure 7 shows a nucleotide sequence (SEQ ID NO:7) of a native sequence PRO4984 cDNA, wherein SEQ ID NO:7 is a clone designated herein as "DNA37155-2651".

35 Figure 8 shows the amino acid sequence (SEQ ID NO:8) derived from the coding sequence of SEQ ID NO:7 shown in Figure 7.

Figure 9 shows a nucleotide sequence (SEQ ID NO:9) of a native sequence PRO4988 cDNA, wherein SEQ ID NO:9 is a clone designated herein as "DNA38269-2654".

Figure 10 shows the amino acid sequence (SEQ ID NO:10) derived from the coding sequence of SEQ ID NO:9 shown in Figure 9.

Figure 11 shows a nucleotide sequence (SEQ ID NO:11) of a native sequence PRO305 cDNA, wherein
5 SEQ ID NO:11 is a clone designated herein as "DNA40619-1220".

Figure 12 shows the amino acid sequence (SEQ ID NO:12) derived from the coding sequence of SEQ ID NO:11 shown in Figure 11.

Figure 13 shows a nucleotide sequence (SEQ ID NO:13) of a native sequence PRO1866 cDNA, wherein
10 SEQ ID NO:13 is a clone designated herein as "DNA44174-2513".

Figure 14 shows the amino acid sequence (SEQ ID NO:14) derived from the coding sequence of SEQ ID NO:13 shown in Figure 13.

Figure 15 shows a nucleotide sequence (SEQ ID NO:15) of a native sequence PRO4996 cDNA, wherein
15 SEQ ID NO:15 is a clone designated herein as "DNA44675-2662".

Figure 16 shows the amino acid sequence (SEQ ID NO:16) derived from the coding sequence of SEQ ID NO:15 shown in Figure 15.

Figure 17 shows a nucleotide sequence (SEQ ID NO:17) of a native sequence PRO4406 cDNA, wherein
20 SEQ ID NO:17 is a clone designated herein as "DNA45408-2615".

Figure 18 shows the amino acid sequence (SEQ ID NO:18) derived from the coding sequence of SEQ ID NO:17 shown in Figure 17.

Figure 19 shows a nucleotide sequence (SEQ ID NO:19) of a native sequence PRO1120 cDNA, wherein
25 SEQ ID NO:19 is a clone designated herein as "DNA48606-1479".

Figure 20 shows the amino acid sequence (SEQ ID NO:20) derived from the coding sequence of SEQ ID NO:19 shown in Figure 19.

Figure 21 shows a nucleotide sequence (SEQ ID NO:21) of a native sequence PRO4990 cDNA, wherein
30 SEQ ID NO:21 is a clone designated herein as "DNA52753-2656".

Figure 22 shows the amino acid sequence (SEQ ID NO:22) derived from the coding sequence of SEQ ID NO:21 shown in Figure 21.

Figure 23 shows a nucleotide sequence (SEQ ID NO:23) of a native sequence PRO738 cDNA, wherein
35 SEQ ID NO:23 is a clone designated herein as "DNA53915-1258".

Figure 24 shows the amino acid sequence (SEQ ID NO:24) derived from the coding sequence of SEQ ID NO:23 shown in Figure 23.

Figure 25 shows a nucleotide sequence (SEQ ID NO:25) of a native sequence PRO3577 cDNA, wherein
SEQ ID NO:25 is a clone designated herein as "DNA53991-2553".

Figure 26 shows the amino acid sequence (SEQ ID NO:26) derived from the coding sequence of SEQ ID NO:25 shown in Figure 25.

Figure 27 shows a nucleotide sequence (SEQ ID NO:27) of a native sequence PRO1879 cDNA, wherein
SEQ ID NO:27 is a clone designated herein as "DNA54009-2517".

Figure 28 shows the amino acid sequence (SEQ ID NO:28) derived from the coding sequence of SEQ ID NO:27 shown in Figure 27.

Figure 29 shows a nucleotide sequence (SEQ ID NO:29) of a native sequence PRO1471 cDNA, wherein SEQ ID NO:29 is a clone designated herein as "DNA56055-1643".

5 Figure 30 shows the amino acid sequence (SEQ ID NO:30) derived from the coding sequence of SEQ ID NO:29 shown in Figure 29.

Figure 31 shows a nucleotide sequence (SEQ ID NO:31) of a native sequence PRO1114 cDNA, wherein SEQ ID NO:31 is a clone designated herein as "DNA57033-1403".

Figure 32 shows the amino acid sequence (SEQ ID NO:32) derived from the coding sequence of SEQ ID NO:31 shown in Figure 31.

10 Figure 33 shows a nucleotide sequence (SEQ ID NO:33) of a native sequence PRO1076 cDNA, wherein SEQ ID NO:33 is a clone designated herein as "DNA57252-1453".

Figure 34 shows the amino acid sequence (SEQ ID NO:34) derived from the coding sequence of SEQ ID NO:33 shown in Figure 33.

15 Figure 35 shows a nucleotide sequence (SEQ ID NO:35) of a native sequence PRO1483 cDNA, wherein SEQ ID NO:35 is a clone designated herein as "DNA58799-1652".

Figure 36 shows the amino acid sequence (SEQ ID NO:36) derived from the coding sequence of SEQ ID NO:35 shown in Figure 35.

Figure 37 shows a nucleotide sequence (SEQ ID NO:37) of a native sequence PRO4985 cDNA, wherein SEQ ID NO:37 is a clone designated herein as "DNA59770-2652".

20 Figure 38 shows the amino acid sequence (SEQ ID NO:38) derived from the coding sequence of SEQ ID NO:37 shown in Figure 37.

Figure 39 shows a nucleotide sequence (SEQ ID NO:39) of a native sequence PRO5000 cDNA, wherein SEQ ID NO:39 is a clone designated herein as "DNA59774-2665".

25 Figure 40 shows the amino acid sequence (SEQ ID NO:40) derived from the coding sequence of SEQ ID NO:39 shown in Figure 39.

Figure 41 shows a nucleotide sequence (SEQ ID NO:41) of a native sequence PRO1881 cDNA, wherein SEQ ID NO:41 is a clone designated herein as "DNA60281-2518".

Figure 42 shows the amino acid sequence (SEQ ID NO:42) derived from the coding sequence of SEQ ID NO:41 shown in Figure 41.

30 Figure 43 shows a nucleotide sequence (SEQ ID NO:43) of a native sequence PRO4314 cDNA, wherein SEQ ID NO:43 is a clone designated herein as "DNA60736-2559".

Figure 44 shows the amino acid sequence (SEQ ID NO:44) derived from the coding sequence of SEQ ID NO:43 shown in Figure 43.

35 Figure 45 shows a nucleotide sequence (SEQ ID NO:45) of a native sequence PRO4987 cDNA, wherein SEQ ID NO:45 is a clone designated herein as "DNA61875-2653".

Figure 46 shows the amino acid sequence (SEQ ID NO:46) derived from the coding sequence of SEQ ID NO:45 shown in Figure 45.

Figure 47 shows a nucleotide sequence (SEQ ID NO:47) of a native sequence PRO4313 cDNA, wherein SEQ ID NO:47 is a clone designated herein as "DNA62312-2558".

Figure 48 shows the amino acid sequence (SEQ ID NO:48) derived from the coding sequence of SEQ ID NO:47 shown in Figure 47.

5 Figure 49 shows a nucleotide sequence (SEQ ID NO:49) of a native sequence PRO4799 cDNA, wherein SEQ ID NO:49 is a clone designated herein as "DNA62849-1604".

Figure 50 shows the amino acid sequence (SEQ ID NO:50) derived from the coding sequence of SEQ ID NO:49 shown in Figure 49.

Figure 51 shows a nucleotide sequence (SEQ ID NO:51) of a native sequence PRO4995 cDNA, wherein SEQ ID NO:51 is a clone designated herein as "DNA66307-2661".

10 Figure 52 shows the amino acid sequence (SEQ ID NO:52) derived from the coding sequence of SEQ ID NO:51 shown in Figure 51.

Figure 53 shows a nucleotide sequence (SEQ ID NO:53) of a native sequence PRO1341 cDNA, wherein SEQ ID NO:53 is a clone designated herein as "DNA66677-2535".

15 Figure 54 shows the amino acid sequence (SEQ ID NO:54) derived from the coding sequence of SEQ ID NO:53 shown in Figure 53.

Figure 55 shows a nucleotide sequence (SEQ ID NO:55) of a native sequence PRO1777 cDNA, wherein SEQ ID NO:55 is a clone designated herein as "DNA71235-1706".

Figure 56 shows the amino acid sequence (SEQ ID NO:56) derived from the coding sequence of SEQ ID NO:55 shown in Figure 55.

20 Figure 57 shows a nucleotide sequence (SEQ ID NO:57) of a native sequence PRO3580 cDNA, wherein SEQ ID NO:57 is a clone designated herein as "DNA71289-2547".

Figure 58 shows the amino acid sequence (SEQ ID NO:58) derived from the coding sequence of SEQ ID NO:57 shown in Figure 57.

25 Figure 59 shows a nucleotide sequence (SEQ ID NO:59) of a native sequence PRO1779 cDNA, wherein SEQ ID NO:59 is a clone designated herein as "DNA73775-1707".

Figure 60 shows the amino acid sequence (SEQ ID NO:60) derived from the coding sequence of SEQ ID NO:59 shown in Figure 59.

Figure 61 shows a nucleotide sequence (SEQ ID NO:61) of a native sequence PRO1754 cDNA, wherein SEQ ID NO:61 is a clone designated herein as "DNA76385-1692".

30 Figure 62 shows the amino acid sequence (SEQ ID NO:62) derived from the coding sequence of SEQ ID NO:61 shown in Figure 61.

Figure 63 shows a nucleotide sequence (SEQ ID NO:63) of a native sequence PRO1906 cDNA, wherein SEQ ID NO:63 is a clone designated herein as "DNA76395-2527".

35 Figure 64 shows the amino acid sequence (SEQ ID NO:64) derived from the coding sequence of SEQ ID NO:63 shown in Figure 63.

Figure 65 shows a nucleotide sequence (SEQ ID NO:65) of a native sequence PRO1870 cDNA, wherein SEQ ID NO:65 is a clone designated herein as "DNA77622-2516".

Figure 66 shows the amino acid sequence (SEQ ID NO:66) derived from the coding sequence of SEQ ID NO:65 shown in Figure 65.

Figure 67 shows a nucleotide sequence (SEQ ID NO:67) of a native sequence PRO4329 cDNA, wherein SEQ ID NO:67 is a clone designated herein as "DNA77629-2573".

5 Figure 68 shows the amino acid sequence (SEQ ID NO:68) derived from the coding sequence of SEQ ID NO:67 shown in Figure 67.

Figure 69 shows a nucleotide sequence (SEQ ID NO:69) of a native sequence PRO4979 cDNA, wherein SEQ ID NO:69 is a clone designated herein as "DNA77645-2648".

Figure 70 shows the amino acid sequence (SEQ ID NO:70) derived from the coding sequence of SEQ ID NO:69 shown in Figure 69.

10 Figure 71 shows a nucleotide sequence (SEQ ID NO:71) of a native sequence PRO1885 cDNA, wherein SEQ ID NO:71 is a clone designated herein as "DNA79302-2521".

Figure 72 shows the amino acid sequence (SEQ ID NO:72) derived from the coding sequence of SEQ ID NO:71 shown in Figure 71.

15 Figure 73 shows a nucleotide sequence (SEQ ID NO:73) of a native sequence PRO1882 cDNA, wherein SEQ ID NO:73 is a clone designated herein as "DNA79865-2519".

Figure 74 shows the amino acid sequence (SEQ ID NO:74) derived from the coding sequence of SEQ ID NO:73 shown in Figure 73.

Figure 75 shows a nucleotide sequence (SEQ ID NO:75) of a native sequence PRO4989 cDNA, wherein SEQ ID NO:75 is a clone designated herein as "DNA80135-2655".

20 Figure 76 shows the amino acid sequence (SEQ ID NO:76) derived from the coding sequence of SEQ ID NO:75 shown in Figure 75.

Figure 77 shows a nucleotide sequence (SEQ ID NO:77) of a native sequence PRO4323 cDNA, wherein SEQ ID NO:77 is a clone designated herein as "DNA80794-2568".

25 Figure 78 shows the amino acid sequence (SEQ ID NO:78) derived from the coding sequence of SEQ ID NO:77 shown in Figure 77.

Figure 79 shows a nucleotide sequence (SEQ ID NO:79) of a native sequence PRO1886 cDNA, wherein SEQ ID NO:79 is a clone designated herein as "DNA80796-2523".

Figure 80 shows the amino acid sequence (SEQ ID NO:80) derived from the coding sequence of SEQ ID NO:79 shown in Figure 79.

30 Figure 81 shows a nucleotide sequence (SEQ ID NO:81) of a native sequence PRO4395 cDNA, wherein SEQ ID NO:81 is a clone designated herein as "DNA80840-2605".

Figure 82 shows the amino acid sequence (SEQ ID NO:82) derived from the coding sequence of SEQ ID NO:81 shown in Figure 81.

35 Figure 83 shows a nucleotide sequence (SEQ ID NO:83) of a native sequence PRO1782 cDNA, wherein SEQ ID NO:83 is a clone designated herein as "DNA80899-2501".

Figure 84 shows the amino acid sequence (SEQ ID NO:84) derived from the coding sequence of SEQ ID NO:83 shown in Figure 83.

Figure 85 shows a nucleotide sequence (SEQ ID NO:85) of a native sequence PRO4338 cDNA, wherein SEQ ID NO:85 is a clone designated herein as "DNA81228-2580".

Figure 86 shows the amino acid sequence (SEQ ID NO:86) derived from the coding sequence of SEQ ID NO:85 shown in Figure 85.

5 Figure 87 shows a nucleotide sequence (SEQ ID NO:87) of a native sequence PRO4341 cDNA, wherein SEQ ID NO:87 is a clone designated herein as "DNA81761-2583".

Figure 88 shows the amino acid sequence (SEQ ID NO:88) derived from the coding sequence of SEQ ID NO:87 shown in Figure 87.

Figure 89 shows a nucleotide sequence (SEQ ID NO:89) of a native sequence PRO5990 cDNA, wherein SEQ ID NO:89 is a clone designated herein as "DNA96042-2682".

10 Figure 90 shows the amino acid sequence (SEQ ID NO:90) derived from the coding sequence of SEQ ID NO:89 shown in Figure 89.

Figure 91 shows a nucleotide sequence (SEQ ID NO:91) of a native sequence PRO3438 cDNA, wherein SEQ ID NO:91 is a clone designated herein as "DNA82364-2538".

15 Figure 92 shows the amino acid sequence (SEQ ID NO:92) derived from the coding sequence of SEQ ID NO:91 shown in Figure 91.

Figure 93 shows a nucleotide sequence (SEQ ID NO:93) of a native sequence PRO4321 cDNA, wherein SEQ ID NO:93 is a clone designated herein as "DNA82424-2566".

Figure 94 shows the amino acid sequence (SEQ ID NO:94) derived from the coding sequence of SEQ ID NO:93 shown in Figure 93.

20 Figure 95 shows a nucleotide sequence (SEQ ID NO:95) of a native sequence PRO4304 cDNA, wherein SEQ ID NO:95 is a clone designated herein as "DNA82430-2557".

Figure 96 shows the amino acid sequence (SEQ ID NO:96) derived from the coding sequence of SEQ ID NO:95 shown in Figure 95.

25 Figure 97 shows a nucleotide sequence (SEQ ID NO:97) of a native sequence PRO1801 cDNA, wherein SEQ ID NO:97 is a clone designated herein as "DNA83500-2506".

Figure 98 shows the amino acid sequence (SEQ ID NO:98) derived from the coding sequence of SEQ ID NO:97 shown in Figure 97.

Figure 99 shows a nucleotide sequence (SEQ ID NO:99) of a native sequence PRO4403 cDNA, wherein SEQ ID NO:99 is a clone designated herein as "DNA83509-2612".

30 Figure 100 shows the amino acid sequence (SEQ ID NO:100) derived from the coding sequence of SEQ ID NO:99 shown in Figure 99.

Figure 101 shows a nucleotide sequence (SEQ ID NO:101) of a native sequence PRO4324 cDNA, wherein SEQ ID NO:101 is a clone designated herein as "DNA83560-2569".

35 Figure 102 shows the amino acid sequence (SEQ ID NO:102) derived from the coding sequence of SEQ ID NO:101 shown in Figure 101.

Figure 103 shows a nucleotide sequence (SEQ ID NO:103) of a native sequence PRO4303 cDNA, wherein SEQ ID NO:103 is a clone designated herein as "DNA84139-2555".

Figure 104 shows the amino acid sequence (SEQ ID NO:104) derived from the coding sequence of SEQ ID NO:103 shown in Figure 103.

Figure 105 shows a nucleotide sequence (SEQ ID NO:105) of a native sequence PRO4305 cDNA, wherein SEQ ID NO:105 is a clone designated herein as "DNA84141-2556".

5 Figure 106 shows the amino acid sequence (SEQ ID NO:106) derived from the coding sequence of SEQ ID NO:105 shown in Figure 105.

Figure 107 shows a nucleotide sequence (SEQ ID NO:107) of a native sequence PRO4404 cDNA, wherein SEQ ID NO:107 is a clone designated herein as "DNA84142-2613".

Figure 108 shows the amino acid sequence (SEQ ID NO:108) derived from the coding sequence of SEQ ID NO:107 shown in Figure 107.

10 Figure 109 shows a nucleotide sequence (SEQ ID NO:109) of a native sequence PRO1884 cDNA, wherein SEQ ID NO:109 is a clone designated herein as "DNA84318-2520".

Figure 110 shows the amino acid sequence (SEQ ID NO:110) derived from the coding sequence of SEQ ID NO:109 shown in Figure 109.

15 Figure 111 shows a nucleotide sequence (SEQ ID NO:111) of a native sequence PRO4349 cDNA, wherein SEQ ID NO:111 is a clone designated herein as "DNA84909-2590".

Figure 112 shows the amino acid sequence (SEQ ID NO:112) derived from the coding sequence of SEQ ID NO:111 shown in Figure 111.

Figure 113 shows a nucleotide sequence (SEQ ID NO:113) of a native sequence PRO4401 cDNA, wherein SEQ ID NO:113 is a clone designated herein as "DNA84912-2610".

20 Figure 114 shows the amino acid sequence (SEQ ID NO:114) derived from the coding sequence of SEQ ID NO:113 shown in Figure 113.

Figure 115 shows a nucleotide sequence (SEQ ID NO:115) of a native sequence PRO1867 cDNA, wherein SEQ ID NO:115 is a clone designated herein as "DNA84925-2514".

25 Figure 116 shows the amino acid sequence (SEQ ID NO:116) derived from the coding sequence of SEQ ID NO:115 shown in Figure 115.

Figure 117 shows a nucleotide sequence (SEQ ID NO:117) of a native sequence PRO4319 cDNA, wherein SEQ ID NO:117 is a clone designated herein as "DNA84928-2564".

Figure 118 shows the amino acid sequence (SEQ ID NO:118) derived from the coding sequence of SEQ ID NO:117 shown in Figure 117.

30 Figure 119 shows a nucleotide sequence (SEQ ID NO:119) of a native sequence PRO4991 cDNA, wherein SEQ ID NO:119 is a clone designated herein as "DNA84932-2657".

Figure 120 shows the amino acid sequence (SEQ ID NO:120) derived from the coding sequence of SEQ ID NO:119 shown in Figure 119.

35 Figure 121 shows a nucleotide sequence (SEQ ID NO:121) of a native sequence PRO4398 cDNA, wherein SEQ ID NO:121 is a clone designated herein as "DNA86592-2607".

Figure 122 shows the amino acid sequence (SEQ ID NO:122) derived from the coding sequence of SEQ ID NO:121 shown in Figure 121.

Figure 123 shows a nucleotide sequence (SEQ ID NO:123) of a native sequence PRO4346 cDNA, wherein SEQ ID NO:123 is a clone designated herein as "DNA86594-2587".

Figure 124 shows the amino acid sequence (SEQ ID NO:124) derived from the coding sequence of SEQ ID NO:123 shown in Figure 123.

5 Figure 125 shows a nucleotide sequence (SEQ ID NO:125) of a native sequence PRO4350 cDNA, wherein SEQ ID NO:125 is a clone designated herein as "DNA86647-2591".

Figure 126 shows the amino acid sequence (SEQ ID NO:126) derived from the coding sequence of SEQ ID NO:125 shown in Figure 125.

Figure 127 shows a nucleotide sequence (SEQ ID NO:127) of a native sequence PRO4318 cDNA, wherein SEQ ID NO:127 is a clone designated herein as "DNA87185-2563".

10 Figure 128 shows the amino acid sequence (SEQ ID NO:128) derived from the coding sequence of SEQ ID NO:127 shown in Figure 127.

Figure 129 shows a nucleotide sequence (SEQ ID NO:129) of a native sequence PRO4340 cDNA, wherein SEQ ID NO:129 is a clone designated herein as "DNA87656-2582".

15 Figure 130 shows the amino acid sequence (SEQ ID NO:130) derived from the coding sequence of SEQ ID NO:129 shown in Figure 129.

Figure 131 shows a nucleotide sequence (SEQ ID NO:131) of a native sequence PRO4400 cDNA, wherein SEQ ID NO:131 is a clone designated herein as "DNA87974-2609".

Figure 132 shows the amino acid sequence (SEQ ID NO:132) derived from the coding sequence of SEQ ID NO:131 shown in Figure 131.

20 Figure 133 shows a nucleotide sequence (SEQ ID NO:133) of a native sequence PRO4320 cDNA, wherein SEQ ID NO:133 is a clone designated herein as "DNA88001-2565".

Figure 134 shows the amino acid sequence (SEQ ID NO:134) derived from the coding sequence of SEQ ID NO:133 shown in Figure 133.

25 Figure 135 shows a nucleotide sequence (SEQ ID NO:135) of a native sequence PRO4409 cDNA, wherein SEQ ID NO:135 is a clone designated herein as "DNA88004-2575".

Figure 136 shows the amino acid sequence (SEQ ID NO:136) derived from the coding sequence of SEQ ID NO:135 shown in Figure 135.

Figure 137 shows a nucleotide sequence (SEQ ID NO:137) of a native sequence PRO4399 cDNA, wherein SEQ ID NO:137 is a clone designated herein as "DNA89220-2608".

30 Figure 138 shows the amino acid sequence (SEQ ID NO:138) derived from the coding sequence of SEQ ID NO:137 shown in Figure 137.

Figure 139 shows a nucleotide sequence (SEQ ID NO:139) of a native sequence PRO4418 cDNA, wherein SEQ ID NO:139 is a clone designated herein as "DNA89947-2618".

35 Figure 140 shows the amino acid sequence (SEQ ID NO:140) derived from the coding sequence of SEQ ID NO:139 shown in Figure 139.

Figure 141 shows a nucleotide sequence (SEQ ID NO:141) of a native sequence PRO4330 cDNA, wherein SEQ ID NO:141 is a clone designated herein as "DNA90842-2574".

Figure 142 shows the amino acid sequence (SEQ ID NO:142) derived from the coding sequence of SEQ ID NO:141 shown in Figure 141.

Figure 143 shows a nucleotide sequence (SEQ ID NO:143) of a native sequence PRO4339 cDNA, wherein SEQ ID NO:143 is a clone designated herein as "DNA91775-2581".

Figure 144 shows the amino acid sequence (SEQ ID NO:144) derived from the coding sequence of SEQ ID NO:143 shown in Figure 143.

Figure 145 shows a nucleotide sequence (SEQ ID NO:145) of a native sequence PRO4326 cDNA, wherein SEQ ID NO:145 is a clone designated herein as "DNA91779-2571".

Figure 146 shows the amino acid sequence (SEQ ID NO:146) derived from the coding sequence of SEQ ID NO:145 shown in Figure 145.

Figure 147 shows a nucleotide sequence (SEQ ID NO:147) of a native sequence PRO6014 cDNA, wherein SEQ ID NO:147 is a clone designated herein as "DNA92217-2697".

Figure 148 shows the amino acid sequence (SEQ ID NO:148) derived from the coding sequence of SEQ ID NO:147 shown in Figure 147.

Figure 149 shows a nucleotide sequence (SEQ ID NO:149) of a native sequence PRO3446 cDNA, wherein SEQ ID NO:149 is a clone designated herein as "DNA92219-2541".

Figure 150 shows the amino acid sequence (SEQ ID NO:150) derived from the coding sequence of SEQ ID NO:149 shown in Figure 149.

Figure 151 shows a nucleotide sequence (SEQ ID NO:151) of a native sequence PRO4322 cDNA, wherein SEQ ID NO:151 is a clone designated herein as "DNA92223-2567".

Figure 152 shows the amino acid sequence (SEQ ID NO:152) derived from the coding sequence of SEQ ID NO:151 shown in Figure 151.

Figure 153 shows a nucleotide sequence (SEQ ID NO:153) of a native sequence PRO4381 cDNA, wherein SEQ ID NO:153 is a clone designated herein as "DNA92225-2603".

Figure 154 shows the amino acid sequence (SEQ ID NO:154) derived from the coding sequence of SEQ ID NO:153 shown in Figure 153.

Figure 155 shows a nucleotide sequence (SEQ ID NO:155) of a native sequence PRO4348 cDNA, wherein SEQ ID NO:155 is a clone designated herein as "DNA92232-2589".

Figure 156 shows the amino acid sequence (SEQ ID NO:156) derived from the coding sequence of SEQ ID NO:155 shown in Figure 155.

Figure 157 shows a nucleotide sequence (SEQ ID NO:157) of a native sequence PRO4371 cDNA, wherein SEQ ID NO:157 is a clone designated herein as "DNA92233-2599".

Figure 158 shows the amino acid sequence (SEQ ID NO:158) derived from the coding sequence of SEQ ID NO:157 shown in Figure 157.

Figure 159 shows a nucleotide sequence (SEQ ID NO:159) of a native sequence PRO3742 cDNA, wherein SEQ ID NO:159 is a clone designated herein as "DNA92243-2549".

Figure 160 shows the amino acid sequence (SEQ ID NO:160) derived from the coding sequence of SEQ ID NO:159 shown in Figure 159.

Figure 161 shows a nucleotide sequence (SEQ ID NO:161) of a native sequence PRO5773 cDNA, wherein SEQ ID NO:161 is a clone designated herein as "DNA92253-2671".

Figure 162 shows the amino acid sequence (SEQ ID NO:162) derived from the coding sequence of SEQ ID NO:161 shown in Figure 161.

5 Figure 163 shows a nucleotide sequence (SEQ ID NO:163) of a native sequence PRO5774 cDNA, wherein SEQ ID NO:163 is a clone designated herein as "DNA92254-2672".

Figure 164 shows the amino acid sequence (SEQ ID NO:164) derived from the coding sequence of SEQ ID NO:163 shown in Figure 163.

Figure 165 shows a nucleotide sequence (SEQ ID NO:165) of a native sequence PRO4343 cDNA, wherein SEQ ID NO:165 is a clone designated herein as "DNA92255-2584".

10 Figure 166 shows the amino acid sequence (SEQ ID NO:166) derived from the coding sequence of SEQ ID NO:165 shown in Figure 165.

Figure 167 shows a nucleotide sequence (SEQ ID NO:167) of a native sequence PRO4325 cDNA, wherein SEQ ID NO:167 is a clone designated herein as "DNA92269-2570".

15 Figure 168 shows the amino acid sequence (SEQ ID NO:168) derived from the coding sequence of SEQ ID NO:167 shown in Figure 167.

Figure 169 shows a nucleotide sequence (SEQ ID NO:169) of a native sequence PRO4347 cDNA, wherein SEQ ID NO:169 is a clone designated herein as "DNA92288-2588".

Figure 170 shows the amino acid sequence (SEQ ID NO:170) derived from the coding sequence of SEQ ID NO:169 shown in Figure 169.

20 Figure 171 shows a nucleotide sequence (SEQ ID NO:171) of a native sequence PRO3743 cDNA, wherein SEQ ID NO:171 is a clone designated herein as "DNA92290-2550".

Figure 172 shows the amino acid sequence (SEQ ID NO:172) derived from the coding sequence of SEQ ID NO:171 shown in Figure 171.

25 Figure 173 shows a nucleotide sequence (SEQ ID NO:173) of a native sequence PRO4426 cDNA, wherein SEQ ID NO:173 is a clone designated herein as "DNA93012-2622".

Figure 174 shows the amino acid sequence (SEQ ID NO:174) derived from the coding sequence of SEQ ID NO:173 shown in Figure 173.

Figure 175 shows a nucleotide sequence (SEQ ID NO:175) of a native sequence PRO4500 cDNA, wherein SEQ ID NO:175 is a clone designated herein as "DNA93020-2642".

30 Figure 176 shows the amino acid sequence (SEQ ID NO:176) derived from the coding sequence of SEQ ID NO:175 shown in Figure 175.

Figure 177 shows a nucleotide sequence (SEQ ID NO:177) of a native sequence PRO4389 cDNA, wherein SEQ ID NO:177 is a clone designated herein as "DNA94830-2604".

35 Figure 178 shows the amino acid sequence (SEQ ID NO:178) derived from the coding sequence of SEQ ID NO:177 shown in Figure 177.

Figure 179 shows a nucleotide sequence (SEQ ID NO:179) of a native sequence PRO4337 cDNA, wherein SEQ ID NO:179 is a clone designated herein as "DNA94833-2579".

Figure 180 shows the amino acid sequence (SEQ ID NO:180) derived from the coding sequence of SEQ ID NO:179 shown in Figure 179.

Figure 181 shows a nucleotide sequence (SEQ ID NO:181) of a native sequence PRO4992 cDNA, wherein SEQ ID NO:181 is a clone designated herein as "DNA94838-2658".

5 Figure 182 shows the amino acid sequence (SEQ ID NO:182) derived from the coding sequence of SEQ ID NO:181 shown in Figure 181.

Figure 183 shows a nucleotide sequence (SEQ ID NO:183) of a native sequence PRO5996 cDNA, wherein SEQ ID NO:183 is a clone designated herein as "DNA94844-2686".

Figure 184 shows the amino acid sequence (SEQ ID NO:184) derived from the coding sequence of SEQ ID NO:183 shown in Figure 183.

10 Figure 185 shows a nucleotide sequence (SEQ ID NO:185) of a native sequence PRO4345 cDNA, wherein SEQ ID NO:185 is a clone designated herein as "DNA94854-2586".

Figure 186 shows the amino acid sequence (SEQ ID NO:186) derived from the coding sequence of SEQ ID NO:185 shown in Figure 185.

15 Figure 187 shows a nucleotide sequence (SEQ ID NO:187) of a native sequence PRO4978 cDNA, wherein SEQ ID NO:187 is a clone designated herein as "DNA95930".

Figure 188 shows the amino acid sequence (SEQ ID NO:188) derived from the coding sequence of SEQ ID NO:187 shown in Figure 187.

Figure 189 shows a nucleotide sequence (SEQ ID NO:189) of a native sequence PRO5780 cDNA, wherein SEQ ID NO:189 is a clone designated herein as "DNA96868-2677".

20 Figure 190 shows the amino acid sequence (SEQ ID NO:190) derived from the coding sequence of SEQ ID NO:189 shown in Figure 189.

Figure 191 shows a nucleotide sequence (SEQ ID NO:191) of a native sequence PRO5992 cDNA, wherein SEQ ID NO:191 is a clone designated herein as "DNA96871-2683".

25 Figure 192 shows the amino acid sequence (SEQ ID NO:192) derived from the coding sequence of SEQ ID NO:191 shown in Figure 191.

Figure 193 shows a nucleotide sequence (SEQ ID NO:193) of a native sequence PRO4428 cDNA, wherein SEQ ID NO:193 is a clone designated herein as "DNA96880-2624".

Figure 194 shows the amino acid sequence (SEQ ID NO:194) derived from the coding sequence of SEQ ID NO:193 shown in Figure 193.

30 Figure 195 shows a nucleotide sequence (SEQ ID NO:195) of a native sequence PRO4994 cDNA, wherein SEQ ID NO:195 is a clone designated herein as "DNA96986-2660".

Figure 196 shows the amino acid sequence (SEQ ID NO:196) derived from the coding sequence of SEQ ID NO:195 shown in Figure 195.

35 Figure 197 shows a nucleotide sequence (SEQ ID NO:197) of a native sequence PRO5995 cDNA, wherein SEQ ID NO:197 is a clone designated herein as "DNA96988-2685".

Figure 198 shows the amino acid sequence (SEQ ID NO:198) derived from the coding sequence of SEQ ID NO:197 shown in Figure 197.

Figure 199 shows a nucleotide sequence (SEQ ID NO:199) of a native sequence PRO6094 cDNA, wherein SEQ ID NO:199 is a clone designated herein as "DNA96995-2709".

Figure 200 shows the amino acid sequence (SEQ ID NO:200) derived from the coding sequence of SEQ ID NO:199 shown in Figure 199.

5 Figure 201 shows a nucleotide sequence (SEQ ID NO:201) of a native sequence PRO4317 cDNA, wherein SEQ ID NO:201 is a clone designated herein as "DNA97004-2562".

Figure 202 shows the amino acid sequence (SEQ ID NO:202) derived from the coding sequence of SEQ ID NO:201 shown in Figure 201.

Figure 203 shows a nucleotide sequence (SEQ ID NO:203) of a native sequence PRO5997 cDNA, wherein SEQ ID NO:203 is a clone designated herein as "DNA97005-2687".

10 Figure 204 shows the amino acid sequence (SEQ ID NO:204) derived from the coding sequence of SEQ ID NO:203 shown in Figure 203.

Figure 205 shows a nucleotide sequence (SEQ ID NO:205) of a native sequence PRO5005 cDNA, wherein SEQ ID NO:205 is a clone designated herein as "DNA97009-2668".

15 Figure 206 shows the amino acid sequence (SEQ ID NO:206) derived from the coding sequence of SEQ ID NO:205 shown in Figure 205.

Figure 207 shows a nucleotide sequence (SEQ ID NO:207) of a native sequence PRO5004 cDNA, wherein SEQ ID NO:207 is a clone designated herein as "DNA97013-2667".

Figure 208 shows the amino acid sequence (SEQ ID NO:208) derived from the coding sequence of SEQ ID NO:207 shown in Figure 207.

20 Figure 209 shows a nucleotide sequence (SEQ ID NO:209) of a native sequence PRO6001 cDNA, wherein SEQ ID NO:209 is a clone designated herein as "DNA98380-2690".

Figure 210 shows the amino acid sequence (SEQ ID NO:210) derived from the coding sequence of SEQ ID NO:209 shown in Figure 209.

25 Figure 211 shows a nucleotide sequence (SEQ ID NO:211) of a native sequence PRO6013 cDNA, wherein SEQ ID NO:211 is a clone designated herein as "DNA98561-2696".

Figure 212 shows the amino acid sequence (SEQ ID NO:212) derived from the coding sequence of SEQ ID NO:211 shown in Figure 211.

Figure 213 shows a nucleotide sequence (SEQ ID NO:213) of a native sequence PRO4502 cDNA, wherein SEQ ID NO:213 is a clone designated herein as "DNA98575-2644".

30 Figure 214 shows the amino acid sequence (SEQ ID NO:214) derived from the coding sequence of SEQ ID NO:213 shown in Figure 213.

Figure 215 shows a nucleotide sequence (SEQ ID NO:215) of a native sequence PRO6007 cDNA, wherein SEQ ID NO:215 is a clone designated herein as "DNA98593-2694".

35 Figure 216 shows the amino acid sequence (SEQ ID NO:216) derived from the coding sequence of SEQ ID NO:215 shown in Figure 215.

Figure 217 shows a nucleotide sequence (SEQ ID NO:217) of a native sequence PRO6028 cDNA, wherein SEQ ID NO:217 is a clone designated herein as "DNA98600-2703".

Figure 218 shows the amino acid sequence (SEQ ID NO:218) derived from the coding sequence of SEQ ID NO:217 shown in Figure 217.

Figure 219 shows a nucleotide sequence (SEQ ID NO:219) of a native sequence PRO100 cDNA, wherein SEQ ID NO:219 is a clone designated herein as "DNA99333".

5 Figure 220 shows the amino acid sequence (SEQ ID NO:220) derived from the coding sequence of SEQ ID NO:219 shown in Figure 219.

Figure 221 shows a nucleotide sequence (SEQ ID NO:221) of a native sequence PRO4327 cDNA, wherein SEQ ID NO:221 is a clone designated herein as "DNA99391-2572".

Figure 222 shows the amino acid sequence (SEQ ID NO:222) derived from the coding sequence of SEQ ID NO:221 shown in Figure 221.

10 Figure 223 shows a nucleotide sequence (SEQ ID NO:223) of a native sequence PRO4315 cDNA, wherein SEQ ID NO:223 is a clone designated herein as "DNA99393-2560".

Figure 224 shows the amino acid sequence (SEQ ID NO:224) derived from the coding sequence of SEQ ID NO:223 shown in Figure 223.

15 Figure 225 shows a nucleotide sequence (SEQ ID NO:225) of a native sequence PRO5993 cDNA, wherein SEQ ID NO:225 is a clone designated herein as "DNA100276-2684".

Figure 226 shows the amino acid sequence (SEQ ID NO:226) derived from the coding sequence of SEQ ID NO:225 shown in Figure 225.

Figure 227 shows a nucleotide sequence (SEQ ID NO:227) of a native sequence PRO4503 cDNA, wherein SEQ ID NO:227 is a clone designated herein as "DNA100312-2645".

20 Figure 228 shows the amino acid sequence (SEQ ID NO:228) derived from the coding sequence of SEQ ID NO:227 shown in Figure 227.

Figure 229 shows a nucleotide sequence (SEQ ID NO:229) of a native sequence PRO4976 cDNA, wherein SEQ ID NO:229 is a clone designated herein as "DNA100902-2646".

25 Figure 230 shows the amino acid sequence (SEQ ID NO:230) derived from the coding sequence of SEQ ID NO:229 shown in Figure 229.

Figure 231 shows a nucleotide sequence (SEQ ID NO:231) of a native sequence PRO5798 cDNA, wherein SEQ ID NO:231 is a clone designated herein as "DNA102899-2679".

Figure 232 shows the amino acid sequence (SEQ ID NO:232) derived from the coding sequence of SEQ ID NO:231 shown in Figure 231.

30 Figure 233 shows a nucleotide sequence (SEQ ID NO:233) of a native sequence PRO6242 cDNA, wherein SEQ ID NO:233 is a clone designated herein as "DNA104875-2720".

Figure 234 shows the amino acid sequence (SEQ ID NO:234) derived from the coding sequence of SEQ ID NO:233 shown in Figure 233.

35 Figure 235 shows a nucleotide sequence (SEQ ID NO:235) of a native sequence PRO6095 cDNA, wherein SEQ ID NO:235 is a clone designated herein as "DNA105680-2710".

Figure 236 shows the amino acid sequence (SEQ ID NO:236) derived from the coding sequence of SEQ ID NO:235 shown in Figure 235.

Figure 237 shows a nucleotide sequence (SEQ ID NO:237) of a native sequence PRO6093 cDNA, wherein SEQ ID NO:237 is a clone designated herein as "DNA105779-2708".

Figure 238 shows the amino acid sequence (SEQ ID NO:238) derived from the coding sequence of SEQ ID NO:237 shown in Figure 237.

5 Figure 239 shows a nucleotide sequence (SEQ ID NO:239) of a native sequence PRO6012 cDNA, wherein SEQ ID NO:239 is a clone designated herein as "DNA105794-2695".

Figure 240 shows the amino acid sequence (SEQ ID NO:240) derived from the coding sequence of SEQ ID NO:239 shown in Figure 239.

Figure 241 shows a nucleotide sequence (SEQ ID NO:241) of a native sequence PRO6027 cDNA, wherein SEQ ID NO:241 is a clone designated herein as "DNA105838-2702".

10 Figure 242 shows the amino acid sequence (SEQ ID NO:242) derived from the coding sequence of SEQ ID NO:241 shown in Figure 241.

Figure 243 shows a nucleotide sequence (SEQ ID NO:243) of a native sequence PRO6181 cDNA, wherein SEQ ID NO:243 is a clone designated herein as "DNA107698-2715".

15 Figure 244 shows the amino acid sequence (SEQ ID NO:244) derived from the coding sequence of SEQ ID NO:243 shown in Figure 243.

Figure 245 shows a nucleotide sequence (SEQ ID NO:245) of a native sequence PRO6097 cDNA, wherein SEQ ID NO:245 is a clone designated herein as "DNA107701-2711".

Figure 246 shows the amino acid sequence (SEQ ID NO:246) derived from the coding sequence of SEQ ID NO:245 shown in Figure 245.

20 Figure 247 shows a nucleotide sequence (SEQ ID NO:247) of a native sequence PRO6090 cDNA, wherein SEQ ID NO:247 is a clone designated herein as "DNA107781-2707".

Figure 248 shows the amino acid sequence (SEQ ID NO:248) derived from the coding sequence of SEQ ID NO:247 shown in Figure 247.

25 Figure 249 shows a nucleotide sequence (SEQ ID NO:249) of a native sequence PRO7171 cDNA, wherein SEQ ID NO:249 is a clone designated herein as "DNA108670-2744".

Figure 250 shows the amino acid sequence (SEQ ID NO:250) derived from the coding sequence of SEQ ID NO:249 shown in Figure 249.

Figure 251 shows a nucleotide sequence (SEQ ID NO:251) of a native sequence PRO6258 cDNA, wherein SEQ ID NO:251 is a clone designated herein as "DNA108688-2725".

30 Figure 252 shows the amino acid sequence (SEQ ID NO:252) derived from the coding sequence of SEQ ID NO:251 shown in Figure 251.

Figure 253 shows a nucleotide sequence (SEQ ID NO:253) of a native sequence PRO9820 cDNA, wherein SEQ ID NO:253 is a clone designated herein as "DNA108769-2765".

35 Figure 254 shows the amino acid sequence (SEQ ID NO:254) derived from the coding sequence of SEQ ID NO:253 shown in Figure 253.

Figure 255 shows a nucleotide sequence (SEQ ID NO:255) of a native sequence PRO6243 cDNA, wherein SEQ ID NO:255 is a clone designated herein as "DNA108935-2721".

Figure 256 shows the amino acid sequence (SEQ ID NO:256) derived from the coding sequence of SEQ ID NO:255 shown in Figure 255.

Figure 257 shows a nucleotide sequence (SEQ ID NO:257) of a native sequence PRO6182 cDNA, wherein SEQ ID NO:257 is a clone designated herein as "DNA110700-2716".

5 Figure 258 shows the amino acid sequence (SEQ ID NO:258) derived from the coding sequence of SEQ ID NO:257 shown in Figure 257.

Figure 259 shows a nucleotide sequence (SEQ ID NO:259) of a native sequence PRO6079 cDNA, wherein SEQ ID NO:259 is a clone designated herein as "DNA111750-2706".

Figure 260 shows the amino acid sequence (SEQ ID NO:260) derived from the coding sequence of SEQ ID NO:259 shown in Figure 259.

10 Figure 261 shows a nucleotide sequence (SEQ ID NO:261) of a native sequence PRO7434 cDNA, wherein SEQ ID NO:261 is a clone designated herein as "DNA123430-2755".

Figure 262 shows the amino acid sequence (SEQ ID NO:262) derived from the coding sequence of SEQ ID NO:261 shown in Figure 261.

15 Figure 263 shows a nucleotide sequence (SEQ ID NO:263) of a native sequence PRO9865 cDNA, wherein SEQ ID NO:263 is a clone designated herein as "DNA125154-2785".

Figure 264 shows the amino acid sequence (SEQ ID NO:264) derived from the coding sequence of SEQ ID NO:263 shown in Figure 263.

Figure 265 shows a nucleotide sequence (SEQ ID NO:265) of a native sequence PRO9828 cDNA, wherein SEQ ID NO:265 is a clone designated herein as "DNA142238-2768".

20 Figure 266 shows the amino acid sequence (SEQ ID NO:266) derived from the coding sequence of SEQ ID NO:265 shown in Figure 265.

Figure 267 shows a nucleotide sequence (SEQ ID NO:267) of a native sequence PRO196 cDNA, wherein SEQ ID NO:267 is a clone designated herein as "DNA22779-1130".

25 Figure 268 shows the amino acid sequence (SEQ ID NO:268) derived from the coding sequence of SEQ ID NO:267 shown in Figure 267.

Figure 269 shows a nucleotide sequence (SEQ ID NO:269) of a native sequence PRO197 cDNA, wherein SEQ ID NO:269 is a clone designated herein as "DNA22780-1078".

Figure 270 shows the amino acid sequence (SEQ ID NO:270) derived from the coding sequence of SEQ ID NO:269 shown in Figure 269.

30 Figure 271 shows a nucleotide sequence (SEQ ID NO:271) of a native sequence PRO195 cDNA, wherein SEQ ID NO:271 is a clone designated herein as "DNA26847-1395".

Figure 272 shows the amino acid sequence (SEQ ID NO:272) derived from the coding sequence of SEQ ID NO:271 shown in Figure 271.

35 Figure 273 shows a nucleotide sequence (SEQ ID NO:273) of a native sequence PRO187 cDNA, wherein SEQ ID NO:273 is a clone designated herein as "DNA27864-1155".

Figure 274 shows the amino acid sequence (SEQ ID NO:274) derived from the coding sequence of SEQ ID NO:273 shown in Figure 273.

Figure 275 shows a nucleotide sequence (SEQ ID NO:275) of a native sequence PRO182 cDNA, wherein SEQ ID NO:275 is a clone designated herein as "DNA27865-1091".

Figure 276 shows the amino acid sequence (SEQ ID NO:276) derived from the coding sequence of SEQ ID NO:275 shown in Figure 275.

5 Figure 277 shows a nucleotide sequence (SEQ ID NO:277) of a native sequence PRO188 cDNA, wherein SEQ ID NO:277 is a clone designated herein as "DNA28497-1130".

Figure 278 shows the amino acid sequence (SEQ ID NO:278) derived from the coding sequence of SEQ ID NO:277 shown in Figure 277.

Figure 279 shows a nucleotide sequence (SEQ ID NO:279) of a native sequence PRO183 cDNA, wherein SEQ ID NO:279 is a clone designated herein as "DNA28498".

10 Figure 280 shows the amino acid sequence (SEQ ID NO:280) derived from the coding sequence of SEQ ID NO:279 shown in Figure 279.

Figure 281 shows a nucleotide sequence (SEQ ID NO:281) of a native sequence PRO184 cDNA, wherein SEQ ID NO:281 is a clone designated herein as "DNA28500".

15 Figure 282 shows the amino acid sequence (SEQ ID NO:282) derived from the coding sequence of SEQ ID NO:281 shown in Figure 281.

Figure 283 shows a nucleotide sequence (SEQ ID NO:283) of a native sequence PRO185 cDNA, wherein SEQ ID NO:283 is a clone designated herein as "DNA28503".

Figure 284 shows the amino acid sequence (SEQ ID NO:284) derived from the coding sequence of SEQ ID NO:283 shown in Figure 283.

20 Figure 285 shows a nucleotide sequence (SEQ ID NO:285) of a native sequence PRO200 cDNA, wherein SEQ ID NO:285 is a clone designated herein as "DNA29101-1122".

Figure 286 shows the amino acid sequence (SEQ ID NO:286) derived from the coding sequence of SEQ ID NO:285 shown in Figure 285.

25 Figure 287 shows a nucleotide sequence (SEQ ID NO:287) of a native sequence PRO202 cDNA, wherein SEQ ID NO:287 is a clone designated herein as "DNA30869".

Figure 288 shows the amino acid sequence (SEQ ID NO:288) derived from the coding sequence of SEQ ID NO:287 shown in Figure 287.

Figure 289 shows a nucleotide sequence (SEQ ID NO:289) of a native sequence PRO214 cDNA, wherein SEQ ID NO:289 is a clone designated herein as "DNA32286-1191".

30 Figure 290 shows the amino acid sequence (SEQ ID NO:290) derived from the coding sequence of SEQ ID NO:289 shown in Figure 289.

Figure 291 shows a nucleotide sequence (SEQ ID NO:291) of a native sequence PRO215 cDNA, wherein SEQ ID NO:291 is a clone designated herein as "DNA32288-1132".

35 Figure 292 shows the amino acid sequence (SEQ ID NO:292) derived from the coding sequence of SEQ ID NO:291 shown in Figure 291.

Figure 293 shows a nucleotide sequence (SEQ ID NO:293) of a native sequence PRO219 cDNA, wherein SEQ ID NO:293 is a clone designated herein as "DNA32290-1164".

Figure 294 shows the amino acid sequence (SEQ ID NO:294) derived from the coding sequence of SEQ ID NO:293 shown in Figure 293.

Figure 295 shows a nucleotide sequence (SEQ ID NO:295) of a native sequence PRO211 cDNA, wherein SEQ ID NO:295 is a clone designated herein as "DNA32292-1131".

5 Figure 296 shows the amino acid sequence (SEQ ID NO:296) derived from the coding sequence of SEQ ID NO:295 shown in Figure 295.

Figure 297 shows a nucleotide sequence (SEQ ID NO:297) of a native sequence PRO220 cDNA, wherein SEQ ID NO:297 is a clone designated herein as "DNA32298-1132".

Figure 298 shows the amino acid sequence (SEQ ID NO:298) derived from the coding sequence of SEQ ID NO:297 shown in Figure 297.

10 Figure 299 shows a nucleotide sequence (SEQ ID NO:299) of a native sequence PRO366 cDNA, wherein SEQ ID NO:299 is a clone designated herein as "DNA33085-1110".

Figure 300 shows the amino acid sequence (SEQ ID NO:300) derived from the coding sequence of SEQ ID NO:299 shown in Figure 299.

15 Figure 301 shows a nucleotide sequence (SEQ ID NO:301) of a native sequence PRO216 cDNA, wherein SEQ ID NO:301 is a clone designated herein as "DNA33087-1158".

Figure 302 shows the amino acid sequence (SEQ ID NO:302) derived from the coding sequence of SEQ ID NO:301 shown in Figure 301.

Figure 303 shows a nucleotide sequence (SEQ ID NO:303) of a native sequence PRO221 cDNA, wherein SEQ ID NO:303 is a clone designated herein as "DNA33089-1132".

20 Figure 304 shows the amino acid sequence (SEQ ID NO:304) derived from the coding sequence of SEQ ID NO:303 shown in Figure 303.

Figure 305 shows a nucleotide sequence (SEQ ID NO:305) of a native sequence PRO228 cDNA, wherein SEQ ID NO:305 is a clone designated herein as "DNA33092-1202".

25 Figure 306 shows the amino acid sequence (SEQ ID NO:306) derived from the coding sequence of SEQ ID NO:305 shown in Figure 305.

Figure 307 shows a nucleotide sequence (SEQ ID NO:307) of a native sequence PRO217 cDNA, wherein SEQ ID NO:307 is a clone designated herein as "DNA33094-1131".

Figure 308 shows the amino acid sequence (SEQ ID NO:308) derived from the coding sequence of SEQ ID NO:307 shown in Figure 307.

30 Figure 309 shows a nucleotide sequence (SEQ ID NO:309) of a native sequence PRO222 cDNA, wherein SEQ ID NO:309 is a clone designated herein as "DNA33107-1135".

Figure 310 shows the amino acid sequence (SEQ ID NO:310) derived from the coding sequence of SEQ ID NO:309 shown in Figure 309.

35 Figure 311 shows a nucleotide sequence (SEQ ID NO:311) of a native sequence PRO224 cDNA, wherein SEQ ID NO:311 is a clone designated herein as "DNA33221-1133".

Figure 312 shows the amino acid sequence (SEQ ID NO:312) derived from the coding sequence of SEQ ID NO:311 shown in Figure 311.

Figure 313 shows a nucleotide sequence (SEQ ID NO:313) of a native sequence PRO230 cDNA, wherein SEQ ID NO:313 is a clone designated herein as "DNA33223-1136".

Figure 314 shows the amino acid sequence (SEQ ID NO:314) derived from the coding sequence of SEQ ID NO:313 shown in Figure 313.

5 Figure 315 shows a nucleotide sequence (SEQ ID NO:315) of a native sequence PRO198 cDNA, wherein SEQ ID NO:315 is a clone designated herein as "DNA33457-1078".

Figure 316 shows the amino acid sequence (SEQ ID NO:316) derived from the coding sequence of SEQ ID NO:315 shown in Figure 315.

Figure 317 shows a nucleotide sequence (SEQ ID NO:317) of a native sequence PRO226 cDNA, wherein SEQ ID NO:317 is a clone designated herein as "DNA33460-1166".

10 Figure 318 shows the amino acid sequence (SEQ ID NO:318) derived from the coding sequence of SEQ ID NO:317 shown in Figure 317.

Figure 319 shows a nucleotide sequence (SEQ ID NO:319) of a native sequence PRO261 cDNA, wherein SEQ ID NO:319 is a clone designated herein as "DNA33473-1176".

15 Figure 320 shows the amino acid sequence (SEQ ID NO:320) derived from the coding sequence of SEQ ID NO:319 shown in Figure 319.

Figure 321 shows a nucleotide sequence (SEQ ID NO:321) of a native sequence PRO242 cDNA, wherein SEQ ID NO:321 is a clone designated herein as "DNA33785-1143".

Figure 322 shows the amino acid sequence (SEQ ID NO:322) derived from the coding sequence of SEQ ID NO:321 shown in Figure 321.

20 Figure 323 shows a nucleotide sequence (SEQ ID NO:323) of a native sequence PRO227 cDNA, wherein SEQ ID NO:323 is a clone designated herein as "DNA33786-1132".

Figure 324 shows the amino acid sequence (SEQ ID NO:324) derived from the coding sequence of SEQ ID NO:323 shown in Figure 323.

25 Figure 325 shows a nucleotide sequence (SEQ ID NO:325) of a native sequence PRO237 cDNA, wherein SEQ ID NO:325 is a clone designated herein as "DNA34353-1428".

Figure 326 shows the amino acid sequence (SEQ ID NO:326) derived from the coding sequence of SEQ ID NO:325 shown in Figure 325.

Figure 327 shows a nucleotide sequence (SEQ ID NO:327) of a native sequence PRO241 cDNA, wherein SEQ ID NO:327 is a clone designated herein as "DNA34392-1170".

30 Figure 328 shows the amino acid sequence (SEQ ID NO:328) derived from the coding sequence of SEQ ID NO:327 shown in Figure 327.

Figure 329 shows a nucleotide sequence (SEQ ID NO:329) of a native sequence PRO231 cDNA, wherein SEQ ID NO:329 is a clone designated herein as "DNA34434-1139".

35 Figure 330 shows the amino acid sequence (SEQ ID NO:330) derived from the coding sequence of SEQ ID NO:329 shown in Figure 329.

Figure 331 shows a nucleotide sequence (SEQ ID NO:331) of a native sequence PRO235 cDNA, wherein SEQ ID NO:331 is a clone designated herein as "DNA35558-1167".